

# Genetic Variation in *CACNA1C* Affects Brain Circuitries Related to Mental Illness

Kristin L. Bigos, PhD; Venkata S. Mattay, MD; Joseph H. Callicott, MD; Richard E. Straub, PhD; Radhakrishna Vakkalanka, PhD; Bhaskar Kolachana, PhD; Thomas M. Hyde, MD; Barbara K. Lipska, PhD; Joel E. Kleinman, MD, PhD; Daniel R. Weinberger, MD

**Context:** The *CACNA1C* gene ( $\alpha$ -1C subunit of the L-type voltage-gated calcium channel) has been identified as a risk gene for bipolar disorder and schizophrenia, but the mechanism of association has not been explored.

**Objective:** To identify the neural system mechanism that explains the genetic association between the *CACNA1C* gene and psychiatric illness using neuroimaging and human brain expression.

**Design:** We used blood oxygenation level-dependent (BOLD) functional magnetic resonance imaging (fMRI) to measure brain activation in circuitries related to bipolar disorder and schizophrenia by comparing *CACNA1C* genotype groups among healthy subjects. We tested the effect of genotype on messenger RNA (mRNA) levels of *CACNA1C* in postmortem human brain. A case-control analysis was used to determine the association of *CACNA1C* genotype with schizophrenia.

**Setting:** National Institutes of Health Clinical Center.

**Patients:** Healthy men and women of white race/ethnicity participated in the fMRI study. Postmortem samples from normal human brains were used for the

brain expression study. Patients with schizophrenia and healthy subjects were used in the case-control analysis.

**Main Outcome Measures:** BOLD fMRI, mRNA levels in postmortem brain samples, and genetic association with schizophrenia.

**Results:** The risk-associated single-nucleotide polymorphism (SNP rs1006737) in *CACNA1C* predicted increased hippocampal activity during emotional processing ( $P = .001$  uncorrected,  $P_{(\text{false recovery rate [FDR]})} = .05$ ,  $z = 3.20$ ) and increased prefrontal activity during executive cognition ( $P = 2.8 \times 10^{-5}$  uncorrected,  $P_{\text{FDR}} = .01$ ,  $z = 4.03$ ). The risk-associated SNP also predicted increased expression of *CACNA1C* mRNA in human brain ( $P = .002$ ). *CACNA1C* was associated with schizophrenia in our case-control sample (odds ratio, 1.77;  $P = .03$ ).

**Conclusions:** The risk-associated SNP in *CACNA1C* maps to circuitries implicated in genetic risk for bipolar disorder and schizophrenia. Its effects in human brain expression implicate a molecular and neural system mechanism for the clinical genetic association.

*Arch Gen Psychiatry.* 2010;67(9):939-945

**Author Affiliations:** Genes, Cognition, and Psychosis Program (Drs Bigos, Mattay, Straub, and Weinberger) and Clinical Brain Disorders Branch (Drs Mattay, Callicott, Straub, Vakkalanka, Kolachana, Hyde, Lipska, Kleinman, and Weinberger), Division of Intramural Research Program, National Institute of Mental Health, National Institutes of Health, Bethesda, Maryland.

SEVERAL RESEARCH GROUPS have performed independent genome-wide association studies<sup>1-3</sup> of bipolar disorder, with little agreement about the most associated loci. However, a comparison of the Wellcome Trust Case Control Consortium and STEP-UCL studies identified *CACNA1C* ( $\alpha$ -1C subunit of the L-type voltage-gated calcium channel) as showing the strongest consistent signal.<sup>3</sup> The most risk-associated single-nucleotide polymorphism (SNP) in this region (rs1006737) across the 2 studies showed association in a separate data set, the so-called ED-DUB-STEP2, as well as in a combined analysis of the 2 data sets (rs1006737,  $P = 7.0 \times 10^{-8}$ ),<sup>4</sup> providing further evidence that *CACNA1C* is a credible susceptibility locus for bipolar disorder.

Because statistical association with clinical diagnosis does not establish biologic significance or identify a mechanism of risk, it is important to extend the statistical evidence with biological data. One approach that has become increasingly informative in translating clinical associations with psychiatric disorders into potential neural mechanisms of risk has been the use of neuroimaging to map gene effects in brain.<sup>5-7</sup> Therefore, we used functional magnetic resonance imaging (fMRI) to test the effects of risk-associated variation in this gene on specific patterns of brain activity that have been associated with mental illness and with increased genetic risk for mental illness. Patients with bipolar disorder have previously been shown to exhibit elevated amygdala<sup>8</sup> and hippocampal<sup>9</sup> activity in response to emo-

**Table 1. Sample Demographics for the Imaging Studies**

Demographic	Genotype			
	Emotional Memory Task			
	<b>GG</b> (n=57)	<b>GA</b> (n=43)	<b>AA</b> (n=16)	<b>All</b> (n=116)
Age, mean (SD), y	29.37 (8.94)	29.47 (9.57)	27.88 (8.02)	29.20 (9.00)
Female sex, No. (%)	30 (52.6)	22 (51.2)	8 (50.0)	60 (51.7)
IQ, mean (SD)	107.8 (7.9)	107.7 (9.6)	107.2 (7.8)	107.6 (8.4)
	Emotional Faces Task			
	<b>GG</b> (n=64)	<b>GA</b> (n=53)	<b>AA</b> (n=14)	<b>All</b> (n=131)
Age, mean (SD), y	29.20 (8.78)	28.98 (8.84)	27.07 (7.98)	28.89 (8.68)
Female sex, No. (%)	34 (53.1)	30 (56.6)	7 (50.0)	71 (54.2)
IQ, mean (SD)	107.9 (8.4)	108.5 (7.6)	106.3 (9.4)	107.9 (8.2)
	Working Memory (N-Back) Task			
	<b>GG</b> (n=146)	<b>GA</b> (n=141)	<b>AA</b> (n=29)	<b>All</b> (n=316)
Age, mean (SD), y	30.92 (9.26)	31.48 (9.55)	30.07 (9.79)	31.09 (9.42)
Female sex, No. (%)	80 (54.8)	74 (52.5)	16 (55.2)	170 (53.8)
IQ, mean (SD)	108.4 (8.9)	109.5 (9.5)	107.6 (9.2)	108.8 (9.2)

tional stimuli. There also is evidence that individuals at increased genetic risk for bipolar disorder show similar patterns of brain activity, suggesting that this may reflect a neural mechanism of genetic risk.<sup>10,11</sup> As a risk gene for bipolar disorder, we hypothesized that healthy subjects who are carriers of the risk-associated allele (A [minor allele]) of *CACNA1C* would have increased amygdala and hippocampal activity in response to emotional stimuli compared with carriers of the common allele (G). Because this gene has recently been associated with risk for schizophrenia, although with less statistical power,<sup>12</sup> we further hypothesized that individuals with this risk-associated allele would show inefficient prefrontal activity during a working memory task, which has been identified as a potential biologic intermediate phenotype related to genetic risk for schizophrenia.<sup>13</sup>

## METHODS

### IMAGING SUBJECTS

Healthy adults participated in an fMRI study in the Clinical Brain Disorders Branch Sibling Study at the National Institute of Mental Health, National Institutes of Health.<sup>14</sup> The study was approved by the National Institute of Mental Health Intramural Program Institutional Review Board. All participants were assessed using the Structured Clinical Interview for *DSM-IV*. All subjects were physically and psychiatrically healthy; specific exclusion criteria have been previously reported.<sup>6</sup> Only white subjects of self-identified European descent were included in this data set to minimize population stratification artifacts. Subject demographics for the imaging study are listed in **Table 1**.

### fMRI TASKS

#### Emotional Memory Task

The emotional memory task involved the encoding and retrieval of aversive scenes,<sup>15</sup> which has been shown to reliably engage the hippocampus in healthy volunteers.<sup>16-18</sup> The scenes, selected from the International Affective Picture System,<sup>19</sup> were presented in a block-designed task with 2 blocks of aversive or

neutral scenes alternating with blocks of resting state for encoding and retrieval blocks. During experimental blocks, 6 scenes of similar valence (neutral or aversive) were presented serially to subjects for 3 seconds each. During resting blocks, participants were asked to attend to a fixation cross presented in the center of the screen for 18 seconds. These fixation blocks were treated as a baseline in the fMRI analyses. During the encoding blocks, subjects were instructed to choose whether the scene depicted an indoor or outdoor scene. During the retrieval blocks, subjects were instructed to select the scenes seen during the encoding session (ie, old) or the scenes not seen during the encoding session (ie, new). In each retrieval block, half of the scenes were old (ie, presented during the encoding session). Each session (encoding or retrieval) consisted of 17 blocks (4 aversive, 4 neutral, and 9 rest conditions). Subjects completed the entire encoding session before beginning the retrieval session after a brief delay (about 2 minutes). The presentation of indoor and outdoor scenes for the encoding session, as well as the presentation of old and new scenes for the retrieval session, was counterbalanced within each block. In addition, the presentation order of aversive and neutral blocks was counterbalanced across subjects. The total imaging time was 5 minutes and 40 seconds for this task. In this study, only the aversive encoding and aversive retrieval tasks were analyzed. There were no significant differences between genotype groups during the aversive retrieval task. BOLD fMRI was performed on a 3-T imaging system (Signa; GE Medical Systems, Milwaukee, Wisconsin) using a gradient-echo echoplanar imaging sequence for all fMRI tasks. Specific parameters for the emotional memory task are the following: 24 axial sections, 4-mm section thickness, 1-mm gap between sections, 2000-millisecond repetition time, 28-millisecond echo time, 24-cm field of view, and 64 × 64-pixel matrix.

#### Emotional Faces Task

The face-matching task is a simple perceptual task, which has previously been shown to robustly engage the amygdala.<sup>5,20,21</sup> The block fMRI paradigm consists of the following 2 experimental conditions: an emotional face-matching condition and a sensorimotor control task. The face-matching task consisted of five 30-second blocks. Blocks 1, 3, and 5 were sensorimotor blocks, and blocks 2 and 4 were emotion blocks. Each sensorimotor and emotion block consisted of six 5-second trials. Each trial involved the presentation of 2 images in the lower panel

and 1 image in the upper panel. In the 6 trials of each sensorimotor block, the 2 lower images were shapes, and the upper panel image was identical to 1 of the shapes in the lower panel. Subjects responded using button presses (left or right) to indicate which lower panel image matches the upper panel image. In the 6 trials of each emotion block, the lower panel consisted of 2 faces, 1 angry and 1 afraid, derived from a standard set of pictures of facial affect.<sup>22</sup> The upper panel consists of 1 of the 2 faces shown in the lower panel. Subjects responded using button presses (left or right) to indicate which lower panel face matches the upper panel face. BOLD fMRI parameters for the emotional faces task are the following: 24 axial sections, 4-mm section thickness, 1-mm gap between sections, 2000-millisecond repetition time, 28-millisecond echo time, 24-cm field of view, and 64 × 64-pixel matrix.

### Working Memory (N-Back) Task

Participants also performed a working memory (N-back) task administered using a block design, with the 2-back working memory condition alternating with a no-back control condition as previously described.<sup>13</sup> During the 0-back control task block, the subject simply responded with the current digit presented (1-4 in a diamond-shaped box). This alternated with the 2-back block in which the subject serially responded with numbers presented 2 previously ("n" = 2). BOLD fMRI parameters for the N-back task are the following: 24 axial sections, 6-mm section thickness, 2000-millisecond repetition time, 28-millisecond echo time, 24-cm field of view, and 64 × 64-pixel matrix.

### IMAGE ANALYSIS

Images were processed as described previously<sup>18</sup> using Statistical Parametric Mapping 5 (SPM5) (<http://www.fil.ion.ucl.ac.uk/spm>). Briefly, images were realigned to the first image of the scan run, spatially normalized into a standard stereotactic space (Montreal Neurological Institute [Quebec City, Quebec, Canada] template) using an affine and nonlinear (4 × 5 × 4 basis functions) transformation, smoothed with a full width at half maximum (FWHM) gaussian filter (8-mm FWHM for emotion tasks and 10-mm for N-back), and ratio normalized to the whole-brain global mean. In the first-level analyses, linear contrasts were computed producing *t* statistical parameter maps at each voxel for emotional tasks. Similarly, *t* statistical parameter maps were produced for the 2-back working memory condition using the 0-back condition as a baseline. These statistical images were entered in a second-level model to identify significant activations within and between genotype groups, thresholded at *P* < .01 uncorrected for the region of interest (ROI) using SPM5. Initially, each of the imaging studies was evaluated using an additive genetic model, with 3 levels of genotype. This was not significant for any of the studies. Therefore, a secondary model (a recessive risk-associated allele model) was tested comparing homozygotes for the risk-associated allele against other genotypes (AA vs GA + GG) using a 2-sample *t* test. Results are presented based on this analysis. For both emotional tasks, the ROI was defined as the amygdala, hippocampus, and parahippocampus using the Wake Forest University, Winston-Salem, North Carolina, Pickatlas.<sup>23</sup> For the N-back task, the ROI was defined as Brodmann areas 9, 10, and 46 using the Pickatlas.<sup>23</sup> No behavioral differences were noted among the 3 genotype groups in the accuracy and reaction times for any of the tasks.

### GENETIC ASSOCIATION COHORT

The cohort used in the genetic association study was the Clinical Brain Disorders Branch–National Institute of Mental Health Sibling Study sample, which consists of subjects collected as

**Table 2. Sample Demographics for the Case-Control Genetic Association Study**

Demographic	Cases (n=282)	Controls (n=440)
Female sex, No. (%)	66 (23.4)	236 (53.6)
Age, mean (SD), y	36.44 (10.54)	33.09 (10.09)
Age at onset, mean (SD), y	21.65 (5.46)	...

**Table 3. Sample Demographics for the Postmortem Human Brain Expression Study**

Demographic	Genotype		
	GG (n=108)	GA (n=116)	AA (n=37)
Age, mean (SD), y	27.35 (22.06)	26.11 (23.39)	31.59 (19.70)
Female sex, No. (%)	36 (33.3)	42 (36.2)	13 (35.1)
Race/ethnicity, No. (%)			
African American	46 (42.6)	72 (62.1)	30 (81.1)
White	62 (57.4)	44 (37.9)	7 (18.9)


part of an ongoing investigation into neurobiological traits related to genetic risk for schizophrenia.<sup>14</sup> Participants were between the ages of 18 and 60 years. To reduce genetic heterogeneity, only white subjects of self-identified European descent were analyzed. Collection details, screening, diagnostic procedures, and exclusion criteria have been previously described.<sup>14</sup> Sample demographics for the genetic association study are listed in **Table 2**.

### GENOTYPING

The *CACNA1C* (OMIM \*114205) SNP rs1006737 was determined in the clinical samples by standard allelic discrimination TaqMan assay that uses the 5' nuclease activity of Taq DNA polymerase to detect a fluorescent reporter signal generated after polymerase chain reaction amplification. The assay cocktail for rs1006737 (Assays on Demand; Applied Biosystems, Foster City, California) was used. Genotype reproducibility was routinely assessed by regenotyping all samples for the selected SNP and was generally greater than 99%. The genotyping completion rate was greater than 95%. Genotypes in all groups were in Hardy-Weinberg equilibrium as determined by an exact test (*P* > .1 for all).

### HUMAN BRAIN EXPRESSION STUDY

Human brain tissue was collected as part of the Clinical Brain Disorders Branch, National Institute of Mental Health brain collection. Details about the collection, screening, and dissection processes for human brain tissues have been described.<sup>24</sup> Expression of mRNA was measured in human dorsolateral prefrontal cortex from prenatal samples (gestational weeks 14-20) and from day of birth through old age using custom microarrays (Illumina; Illumina, Inc, San Diego, California). Sample demographics for the postmortem human brain tissue study are listed in **Table 3**. Microarray chips were generated in the National Human Genome Research Institute Microarray Core Facility from 44 544 probes (70mer) obtained from the Illumina Oligoset Human Exonic Evidence Based Oligonucleotide arrays (<http://www.microarray.org/sfgt/heebo.do>). The RNA samples (500 ng) from the Clinical Brain Disorders Branch brain series



**Figure 1.** Risk-associated allele homozygotes (AA) have a trend for greater hippocampal activity during encoding of aversive images (19 voxels [peak voxel 22, -11, -11] for the right hippocampus; 28 voxels [peak voxel -19, -15, -15] for the left hippocampus).

across the lifespan collection were amplified and labeled using fluorescent dye (Cy5). Samples were each hybridized to microarrays simultaneously with a reference standard (labeled with Cy3) consisting of a pool of RNA from many brain tissue samples. The data were normalized using the loess method from the Limma R package (<http://www.bioconductor.org/packages/2.6/bioc/html/limma.html>),<sup>25</sup> and data outside 3 SDs were treated as outliers. Human fetal tissue was obtained from the Eunice Kennedy Shriver National Institute of Child Health and Human Development Brain and Tissue Bank for Developmental Disorders at the University of Maryland, Baltimore. Genotyping of these samples was performed using SNP chips (Illumina 1M Infinium, Illumina, Inc). The rs1006737 is not found on this SNP platform, so a proxy SNP was selected for genetic analysis based on linkage disequilibrium in the sample (HapMap-CEU [[http://www.ncbi.nlm.nih.gov/SNP/snp\\_viewTable.cgi?pop=1409](http://www.ncbi.nlm.nih.gov/SNP/snp_viewTable.cgi?pop=1409)]). Subjects were genotyped for CACNA1C rs2159100, which is in allelic identity with rs1006737 ( $R^2 = 1.0$ ).<sup>26</sup> Because the postmortem data set contained a larger percentage of minor allele carriers than the clinical samples, only an additive genetic model was tested using linear regression analysis to determine the effect of genotype on CACNA1C expression based on 3 genotype groups.

## RESULTS

For each imaging task, subjects in each genotype group were matched for age, sex, IQ, and task performance, thus isolating the effect of genotype on brain information processing not confounded by general brain function parameters or by task performance. The first 2 tasks focused on the engagement of medial temporal lobe structures implicated in emotional processing and associated with mood disorders and increased genetic risk for mood disorders.

### EMOTIONAL IMAGING TASKS

The first task involved the encoding and retrieval of aversive stimuli. For this emotional memory task, images from 116 subjects (57 GG genotype, 43 GA, and 16 AA) were


used in an ROI analysis of the amygdala and hippocampus. Initial testing of an additive genetic model yielded nonsignificant results. Testing of the recessive genetic model showed an effect of genotype on the engagement of the hippocampus. Homozygotes for the risk-associated allele (AA) had greater bilateral hippocampal activity during the encoding of aversive images compared with carriers of the common allele ( $P = .001$  uncorrected,  $P_{(\text{false recovery rate [FDR]})} = .05$ ,  $z = 3.20$  for the right hippocampus;  $P = .003$  uncorrected,  $P_{\text{FDR}} = .05$ ,  $z = 2.77$  for the left hippocampus) (**Figure 1**). Although the uncorrected  $P$  values were significant after correcting for the 2 different genetic models tested, the FDR-corrected results remained significant only at the trend level.

The second task involved matching of emotional faces (ie, angry or fearful), which has been previously shown to robustly activate the amygdala and the hippocampus.<sup>5</sup> For this emotional faces task, images from 131 subjects (64 GG genotype, 53 GA, and 14 AA) were used in an ROI analysis of the amygdala and hippocampus. There were no regions in the amygdala or hippocampus that survived a threshold of  $P < .01$  uncorrected; however, when thresholded at  $P < .05$ , risk-associated allele homozygotes (AA) had slightly greater right amygdala activity compared with carriers of the common allele ( $P = .02$  uncorrected,  $P_{\text{FDR}} = .21$ ,  $z = 2.17$ ). Although this activation was not significant by correction for multiple voxels within the region, it was similar to the pattern of hippocampal activation during encoding of aversive scenes.

### WORKING MEMORY TASK

We also examined the effect of CACNA1C genotype on a task not related to limbic processing of emotion and not previously associated with mood disorders but linked with schizophrenia and prefrontal cognitive processing. If this gene is associated more strongly with mood disorders than with schizophrenia because of a primary effect on emotional circuitry, then a task targeting emotionally neutral cognitive processing related to prefrontal cortex should show a less robust effect. We chose the N-back, a working memory task that robustly engages prefrontal cortical circuitry and has been especially useful in characterizing neural mechanisms of genes associated with schizophrenia.<sup>6,27</sup> Images from 316 subjects (146 GG genotype, 141 GA, and 29 AA), comparing 2-back with 0-back, were analyzed using an ROI for Brodmann areas 9, 10, and 46. Again, analysis under an additive genetic model was nonsignificant. However, testing of the recessive genetic model showed a significant effect of genotype on the efficiency of prefrontal cortex. Homozygotes for the risk-associated allele (AA) had greater activity in prefrontal cortex ( $P = 2.8 \times 10^{-5}$  uncorrected,  $P_{\text{FDR}} = .01$ ,  $z = 4.03$  for the first cluster;  $P = 5.67 \times 10^{-5}$  uncorrected,  $P_{\text{FDR}} = .01$ ,  $z = 3.86$  for the second cluster) (**Figure 2**), a pattern of inefficient engagement previously found in association with several putative schizophrenia susceptibility genotypes.<sup>6,27</sup> Even using a whole-brain analysis, these 2 clusters survived the threshold of  $P < .001$  and 10 contiguous voxels, which suggests that the effect of the CACNA1C gene on this working memory task is specific to prefrontal cortex. These results also remained significant after correcting for the 2





**Figure 2.** Risk-associated allele homozygotes (AA) have prefrontal cortical inefficiency (greater activity) during the N-back working memory task ( $P < .001$  for whole-brain analysis; 88 voxels [peak voxel 54, 12, 39] for the first cluster; 20 voxels [peak voxel 33, 45, 15] for the second cluster). Red areas are significant voxels.


different genetic models tested. Although the significance level seemed greater in prefrontal cortex during this working memory task compared with the hippocampal response in the emotional memory task, this was likely due to a difference in sample size, as the effect sizes are similar (0.76 for emotional memory and 0.78 for working memory).

#### CLINICAL GENETIC ASSOCIATION WITH SCHIZOPHRENIA

The finding of an intermediate brain phenotype associated with genetic risk for schizophrenia<sup>10</sup> suggests that *CACNA1C* might also show an association with the clinical diagnosis of schizophrenia, and indeed this has recently been reported, including a study<sup>12</sup> with the same SNP as that reported herein (rs1006737,  $P = .03$ ). Genome-wide association results of the International Schizophrenia Consortium showed an association of schizophrenia with *CACNA1C*, although not to the same SNP (rs2238090,  $P = 7.7 \times 10^{-6}$ ).<sup>28</sup> We examined *CACNA1C* rs1006737 in a case-control analysis (282 cases and 440 controls) and found a nominal association ( $P = .03$ ) with schizophrenia (odds ratio for risk-associated allele homozygotes, 1.77; 95% confidence interval, 1.07-2.91).

#### ASSOCIATION WITH EXPRESSION OF *CACNA1C* mRNA IN HUMAN BRAIN

*CACNA1C* is highly expressed in heart and in brain. To understand the molecular mechanism underlying the clinical association and the functional differences in brain circuitry, we tested the effect of genetic variation in *CACNA1C* on mRNA expression in a large cohort of human post-mortem brain samples. The proxy SNP (rs2159100) for rs1006737 ( $R^2 = 1.0$ ) showed a significant effect on *CACNA1C* expression; carriers of the risk-associated genotype (AA) had highest expression, with heterozygotes (GA) having intermediate expression and the common allele carriers (GG) having the lowest expression ( $P = .002$  for linear regression analysis) (**Figure 3** and eTable; available



**Figure 3.** Risk-associated allele homozygotes (AA) have greater expression of *CACNA1C* than heterozygotes and common allele homozygotes (probe 28032) (rs2159100,  $P = .002$  for linear regression analysis).

at <http://www.archgenpsychiatry.com>). Because of the small size of the postmortem sample, we also generated an empirical  $P$  value by permutation analysis, which involved randomizing the genotypes and level of expression and 100 000 repetitions of the regression analysis, yielding an empirical  $P = .002$ . Expression of the probe did not significantly change across age, and genotype groups did not differ in mean age; therefore, age was not used as a covariate. There was a difference in expression between persons of white vs African American race/ethnicity, which is likely due to a difference in minor allele frequency (26% vs 45%, respectively). However, there was not a race/ethnicity  $\times$  genotype interaction, implicating genotype as an independent predictor of expression. An analysis using race/ethnicity as a covariate showed a significant effect of genotype on expression ( $F = 3.275$ ,  $P = .04$ ).

#### COMMENT

We found that genetic variation, previously implicated as a risk factor for bipolar disorder and for schizophrenia, shows effects on brain functions related to medio-temporal emotional processing and prefrontal cortical working memory processing that have been associated with risk for bipolar disorder and for schizophrenia, respectively. Our results suggest that the pleiotropic effects of the risk-associated genotype on these diverse brain circuits parallel the diagnostic nonspecificity of the clinical associations and may reflect the underlying neural system mechanisms involved. Genetic variation in voltage-gated calcium channel genes has been associated with several other complex multigenic neuropsychiatric disorders, including autism,<sup>29</sup> epilepsy and migraine,<sup>30</sup> and schizophrenia.<sup>28,31</sup> In addition, a missense mutation in *CACNA1C* results in Timothy syndrome, which is characterized by multiorgan dysfunction, including cardiac arrhythmias and cognitive abnormalities.<sup>32</sup>

Data in this study suggest that calcium channel dysfunction may contribute in part to the genetic etiology of bipolar disorder and schizophrenia through alterations in the functional activity of brain circuitries implicated in both conditions. This is analogous to other genes (eg, *COMT*,<sup>6</sup>

GRM3,<sup>27</sup> BDNF,<sup>18</sup> and DISC1<sup>33</sup>) that have been associated with both diagnoses and both patterns of neural circuitry effects. Investigations of the N-back working memory task have shown that patients with schizophrenia and their healthy siblings have increased prefrontal cortical activity for a given level of performance,<sup>13</sup> suggesting that inefficiency in this circuitry is heritable and a good intermediate phenotype related to genetic risk for schizophrenia. Analogous studies<sup>8,9</sup> have been performed among patients with bipolar disorder using neuroimaging tasks that target mood circuitry in the temporal lobe. Other studies have targeted serotonin signaling genes and have found that healthy subjects who are carriers of the short allele of the serotonin transporter, the target of drugs that treat the mood symptoms of bipolar disorder, have higher ratings of anxiety and depression<sup>34</sup> and have greater activation of the limbic circuitry,<sup>5,20</sup> similar to these data. Although we found stronger statistical evidence of association with prefrontal processing, this was likely an artifact of the reduced power of the smaller sample in the emotional processing tasks, as the effect sizes were similar.

The molecular mechanism of genetic risk seems to relate at least in part to regulation of gene expression. Carriers of the risk-associated allele had increased levels of CACNA1C mRNA. It may be that a specific transcript of CACNA1C that is measured by the oligonucleotide probe is involved in the function of the brain regions measured by fMRI in this study. Calcium channels are involved in various aspects of neuronal development and in the establishment of maintenance of connectivity during development and throughout adulthood<sup>35</sup>; therefore, alterations in gene expression may affect brain structure, which could also affect brain function. A recent study<sup>36</sup> reported that variation in CACNA1C results in alterations in cerebral gray matter. Total gray matter was highest in AA carriers of the risk-associated SNP (rs1006737) compared with GA and GA carriers. However, it is unclear whether and how this finding relates to the genetic association with psychiatric illness.

Our data add to the growing literature that genes weakly associated with psychiatric diseases show stronger effects at the level of brain processing of emotional and cognitive information. This has been shown for several other genes that have been found to be positively but weakly associated with clinical diagnosis in large population studies, only to show strong effects in imaging among much smaller samples. For example, a recent study<sup>37</sup> of the ZNF804A gene, which showed significant genome-wide association with schizophrenia ( $P < 10^{-8}$ ) in a sample of more than 23 000, also showed strong association ( $P = .006$ ) with an imaging phenotype in a much smaller sample ( $n = 115$ ) of healthy subjects. We can explore this question of relative statistical power directly in our data compared with the results of the large-scale clinical study<sup>4</sup> of CACNA1C that showed genome-wide significant association ( $P < 10^{-8}$ ) in a sample of 10 596 total subjects. Using the imaging data from our N-back study, in which  $P = 2.8 \times 10^{-5}$  for a sample size of 316 subjects, if we increase the sample size to 10 000 to approximate the sample size in the previous genome-wide association studies and hold the effect size constant, our  $P$ -value drops to  $P = 4.87 \times 10^{-9}$ . By the same token, using our postmortem expression data, in

which  $P = .002$  for a sample size of 261, if the sample size was increased to 10 000, the  $P$ -value would be  $P = 1.24 \times 10^{-4}$ . It should come as no surprise that in principle our findings of associations with quantitative biological measures related to clinical illness and to genetic risk for illness show greater effect sizes than association with clinical diagnosis for the following 2 reasons: (1) a quantitative trait phenotype has greater statistical power in general compared with a categorical variable and (2) the measures we studied in brain are likely closer to the neurobiological function of the gene and its effect on illness than is the clinical diagnosis. Analogous findings have been reported in other areas of complex medical genetics; for example, genes that show weak association with complex syndromes such as hypertension and cardiovascular disease show much stronger statistical association with quantitative biological traits related to these syndromes (ie, sodium homeostasis<sup>38</sup> and lipid metabolism,<sup>39</sup> respectively).

The association of CACNA1C with brain-related quantitative phenotypes has potential clinical implications. Our demonstration of increased expression of the CACNA1C transcript suggests that, if this translates into increased calcium channel activity, calcium channel inhibitors may have clinical value in treating these disorders. Indeed, anecdotal reports and results of small clinical trials suggest benefit of these agents for some patients,<sup>40,41</sup> but the data have been inconsistent and limited. Genotype or brain imaging-based phenotypes might be considered as individual predictors of response to these agents in future trials. Although further studies are necessary to fully characterize the mechanism by which alterations in CACNA1C expression result in brain function changes, this study identifies a potential mechanism of risk for bipolar disorder and schizophrenia.

**Submitted for Publication:** August 25, 2009; final revision received January 25, 2010; accepted March 5, 2010.

**Correspondence:** Daniel R. Weinberger, MD, Genes, Cognition, and Psychosis Program, Division of Intramural Research Program, National Institute of Mental Health, National Institutes of Health, 10 Center Dr, MSC 1379, Bethesda, MD 20892 (weinberd@mail.nih.gov).

**Financial Disclosure:** None reported.

**Funding/Support:** This research was supported by the Intramural Research Program of the National Institute of Mental Health, National Institutes of Health.

**Online-Only Material:** The eTable is available at <http://www.archgenpsychiatry.com>.

## REFERENCES

1. Baum AE, Akula N, Cabanero M, Cardona I, Corona W, Klemens B, Schulze TG, Cichon S, Rietschel M, Nöthen MM, Georgi A, Schumacher J, Schwarz M, Abou Jamra R, Höfels S, Propping P, Satagopan J, Detera-Wadleigh SD, Hardy J, McMahon FJ. A genome-wide association study implicates diacylglycerol kinase eta (DGKH) and several other genes in the etiology of bipolar disorder. *Mol Psychiatry*. 2008;13(2):197-207.
2. Wellcome Trust Case Control Consortium. Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature*. 2007;447(7145):661-678.
3. Sklar P, Smoller JW, Fan J, Ferreira MA, Perlis RH, Chambert K, Nimgaonkar VL, McQueen MB, Faraone SV, Kirby A, de Bakker PI, Ogdie MN, Thase ME, Sachs GS, Todd-Brown K, Gabriel SB, Sougnez C, Gates C, Blumenstiel B, Defelice M,

- Ardlie KG, Franklin J, Muir WJ, McGhee KA, MacIntyre DJ, McLean A, VanBeck M, McQuillin A, Bass NJ, Robinson M, Lawrence J, Anjorin A, Curtis D, Scolnick EM, Daly MJ, Blackwood DH, Gurling HM, Purcell SM. Whole-genome association study of bipolar disorder. *Mol Psychiatry*. 2008;13(6):558-569.
4. Ferreira MAR, O'Donovan MC, Meng YA, Jones IR, Ruderfer DM, Jones L, Fan J, Kirov G, Perlis RH, Green EK, Smoller JW, Grozeva D, Stone J, Nikolov I, Chambert K, Hamshere ML, Nimgaonkar VL, Moskvina V, Thase ME, Caesar S, Sachs GS, Franklin J, Gordon-Smith K, Ardlie KG, Gabriel SB, Fraser C, Blumenstiel B, Defelice M, Breen G, Gill M, Morris DW, Elkin A, Muir WJ, McGhee KA, Williamson R, MacIntyre DJ, MacLean AW, St CD, Robinson M, Van Beck M, Pereira AC, Kandaswamy R, McQuillin A, Collier DA, Bass NJ, Young AH, Lawrence J, Ferrier IN, Anjorin A, Farmer A, Curtis D, Scolnick EM, McGuffin P, Daly MJ, Corvin AP, Holmans PA, Blackwood DH, Gurling HM, Owen MJ, Purcell SM, Sklar P, Craddock N; Wellcome Trust Case Control Consortium. Collaborative genome-wide association analysis supports a role for *ANKK3* and *CACNA1C* in bipolar disorder. *Nat Genet*. 2008;40(9):1056-1058.
  5. Hariri AR, Mattay VS, Tessitore A, Kolachana B, Fera F, Goldman D, Egan MF, Weinberger DR. Serotonin transporter genetic variation and the response of the human amygdala. *Science*. 2002;297(5580):400-403.
  6. Egan MF, Goldberg TE, Kolachana BS, Callicott JH, Mazzanti CM, Straub RE, Goldman D, Weinberger DR. Effect of *COMT* Val<sup>108/158</sup> Met genotype on frontal lobe function and risk for schizophrenia. *Proc Natl Acad Sci U S A*. 2001;98(12):6917-6922.
  7. Meyer-Lindenberg A, Weinberger DR. Intermediate phenotypes and genetic mechanisms of psychiatric disorders. *Nat Rev Neurosci*. 2006;7(10):818-827.
  8. Almeida JR, Versace A, Hassel S, Kupfer DJ, Phillips ML. Elevated amygdala activity to sad facial expressions: a state marker of bipolar but not unipolar depression. *Biol Psychiatry*. 2010;67(5):414-421.
  9. Whalley HC, McKirdy J, Romaniuk L, Sussmann J, Johnstone EC, Wan HI, McIntosh AM, Lawrie SM, Hall J. Functional imaging of emotional memory in bipolar disorder and schizophrenia. *Bipolar Disord*. 2009;11(8):840-856.
  10. Krüger S, Alda M, Young LT, Goldapple K, Parikh S, Mayberg HS. Risk and resilience markers in bipolar disorder: brain responses to emotional challenge in bipolar patients and their healthy siblings. *Am J Psychiatry*. 2006;163(2):257-264.
  11. Haldane M, Kempton M, Frangou S. Ventral prefrontal function mediates resilience to bipolar disorder: an fMRI study of BD patients and their unaffected siblings [abstract]. *Eur Psychiatry*. 2009;24(suppl 1):S916.
  12. Green EK, Grozeva D, Jones I, Jones L, Kirov G, Caesar S, Gordon-Smith K, Fraser C, Forty L, Russell E, Hamshere ML, Moskvina V, Nikolov I, Farmer A, McGuffin P, Holmans PA, Owen MJ, O'Donovan MC, Craddock N; Wellcome Trust Case Control Consortium 5. The bipolar disorder risk allele at *CACNA1C* also confers risk of recurrent major depression and of schizophrenia [published online ahead of print July 21, 2009]. *Mol Psychiatry*. doi:10.1038/mp.2009.49.
  13. Callicott JH, Egan MF, Mattay VS, Bertolino A, Bone AD, Verchinski B, Weinberger DR. Abnormal fMRI response of the dorsolateral prefrontal cortex in cognitively intact siblings of patients with schizophrenia. *Am J Psychiatry*. 2003;160(4):709-719.
  14. Egan MF, Goldberg TE, Gscheidle T, Weirich M, Bigelow LB, Weinberger DR. Relative risk of attention deficits in siblings of patients with schizophrenia. *Am J Psychiatry*. 2000;157(8):1309-1316.
  15. Murty VP, Sambataro F, Das S, Tan HY, Callicott JH, Goldberg TE, Meyer-Lindenberg A, Weinberger DR, Mattay VS. Age-related alterations in simple declarative memory and the effect of negative stimulus valence. *J Cogn Neurosci*. 2009;21(10):1920-1933.
  16. Bertolino A, Rubino V, Sambataro F, Blasi G, Latorre V, Fazio L, Caforio G, Petruzella V, Kolachana B, Hariri A, Meyer-Lindenberg A, Nardini M, Weinberger DR, Scarabino T. Prefrontal-hippocampal coupling during memory processing is modulated by *COMT* val158met genotype. *Biol Psychiatry*. 2006;60(11):1250-1258.
  17. Meyer-Lindenberg A, Buckholz JW, Kolachana B, R Hariri A, Pezawas L, Blasi G, Wabnitz A, Honea R, Verchinski B, Callicott JH, Egan M, Mattay V, Weinberger DR. Neural mechanisms of genetic risk for impulsivity and violence in humans. *Proc Natl Acad Sci U S A*. 2006;103(16):6269-6274.
  18. Hariri AR, Goldberg TE, Mattay VS, Kolachana BS, Callicott JH, Egan MF, Weinberger DR. Brain-derived neurotrophic factor val<sup>66</sup>met polymorphism affects human memory-related hippocampal activity and predicts memory performance. *J Neurosci*. 2003;23(17):6690-6694.
  19. Lang PJ, Bradley MM, Cuthbert BN. *International Affective Picture System (IAPS): Affective Ratings of Pictures and Instruction Manual*. Gainesville: University of Florida; 2005. Technical report A-6.
  20. Hariri AR, Drabant EM, Munoz KE, Kolachana BS, Mattay VS, Egan MF, Weinberger DR. A susceptibility gene for affective disorders and the response of the human amygdala. *Arch Gen Psychiatry*. 2005;62(2):146-152.
  21. Rasetti R, Mattay VS, Wiedholz LM, Kolachana BS, Hariri AR, Callicott JH, Meyer-Lindenberg A, Weinberger DR. Evidence that altered amygdala activity in schizophrenia is related to clinical state and not genetic risk. *Am J Psychiatry*. 2009;166(2):216-225.
  22. Ekman P, Friesen WV. *Pictures of Facial Affect*. Palo Alto, CA: Consulting Psychologists Press; 1976.
  23. Maldjian JA, Laurienti PJ, Kraft RA, Burdette JH. An automated method for neuroanatomic and cytoarchitectonic atlas-based interrogation of fMRI data sets. *Neuroimage*. 2003;19(3):1233-1239.
  24. Lipska BK, Peters T, Hyde TM, Halim N, Horowitz C, Mitkus S, Weickert CS, Matsumoto M, Sawa A, Straub RE, Vakkalanka R, Herman MM, Weinberger DR, Kleinman JE. Expression of DISC1 binding partners is reduced in schizophrenia and associated with DISC1 SNPs. *Hum Mol Genet*. 2006;15(8):1245-1258.
  25. Smyth GK. Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Stat Appl Genet Mol Biol*. 2004;3(1):e3.
  26. Johnson AD, Handsaker RE, Pulit SL, Nizzari MM, O'Donnell CJ, de Bakker PIW. SNAP: a web-based tool for identification and annotation of proxy SNPs using HapMap. *Bioinformatics*. 2008;24(24):2938-2939.
  27. Tan HY, Chen Q, Sust S, Buckholz JW, Meyers JD, Egan MF, Mattay VS, Meyer-Lindenberg A, Weinberger DR, Callicott JH. Epistasis between catechol-O-methyltransferase and type II metabotropic glutamate receptor 3 genes on working memory brain function. *Proc Natl Acad Sci U S A*. 2007;104(30):12536-12541.
  28. Purcell SM, Wray NR, Stone JL, Visscher PM, O'Donovan MC, Sullivan PF, Sklar P; International Schizophrenia Consortium. Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. *Nature*. 2009;460(7256):748-752.
  29. Wang K, Zhang H, Ma D, Bucan M, Glessner JT, Abrahams BS, Salyakina D, Imielinski M, Bradfield JP, Sleiman PM, Kim CE, Hou C, Frackelton E, Chiavacci R, Takahashi N, Sakurai T, Rappaport E, Lajonchere CM, Munson J, Estes A, Korvatska O, Piven J, Sonnenblick LI, Alvarez Retuerto AI, Herman EI, Dong H, Hutman T, Sigman M, Ozonoff S, Klin A, Owley T, Sweeney JA, Brune CW, Cantor RM, Bernier R, Gilbert JR, Cuccaro ML, McMahon WM, Miller J, State MW, Wassink TH, Coon H, Levy SE, Schultz RT, Nurnberger Jr, Haines JL, Sutcliffe JS, Cook EH, Minshew NJ, Buxbaum JD, Dawson G, Grant SF, Geschwind DH, Pericak-Vance MA, Schellenberg GD, Hakonarson H. Common genetic variants on 5p14.1 associate with autism spectrum disorders. *Nature*. 2009;459(7246):528-533.
  30. Gargus JJ. Genetic calcium signaling abnormalities in the central nervous system: seizures, migraine, and autism. *Ann N Y Acad Sci*. 2009;1151:133-156.
  31. Wei J, Hemmings GP. A further study of a possible locus for schizophrenia on the X chromosome. *Biochem Biophys Res Commun*. 2006;344(4):1241-1245.
  32. Splawski I, Timothy KW, Sharpe LM, Decher N, Kumar P, Bloise R, Napolitano C, Schwartz PJ, Joseph RM, Condouris K, Tager-Flusberg H, Priori SG, Sanguinetti MC, Keating MT. Ca<sub>v</sub>1.2 calcium channel dysfunction causes a multisystem disorder including arrhythmia and autism. *Cell*. 2004;119(1):19-31.
  33. Callicott JH, Straub RE, Pezawas L, Egan MF, Mattay VS, Hariri AR, Verchinski BA, Meyer-Lindenberg A, Balkissoon R, Kolachana B, Goldberg TE, Weinberger DR. Variation in *DISC1* affects hippocampal structure and function and increases risk for schizophrenia. *Proc Natl Acad Sci U S A*. 2005;102(24):8627-8632.
  34. Bigos KL, Hariri AR. Neuroimaging: technologies at the interface of genes, brain, and behavior. *Neuroimaging Clin N Am*. 2007;17(4):459-467, viii.
  35. Spitzer NC. Electrical activity in early neuronal development. *Nature*. 2006;444(7120):707-712.
  36. Kempton MJ, Ruberto G, Vassos E, Tatarelli R, Girardi P, Collier D, Frangou S. Effects of the *CACNA1C* risk allele for bipolar disorder on cerebral gray matter volume in healthy individuals [letter]. *Am J Psychiatry*. 2009;166(12):1413-1414.
  37. Esslinger C, Walter H, Kirsch P, Erk S, Schnell K, Arnold C, Haddad L, Mier D, Opitz von Boberfeld C, Raab K, Witt SH, Rietschel M, Cichon S, Meyer-Lindenberg A. Neural mechanisms of a genome-wide supported psychosis variant. *Science*. 2009;324(5927):605.
  38. Newton-Cheh C, Larson MG, Vasas RS, Levy D, Bloch KD, Surati A, Guiducci C, Kathiresan S, Benjamin EJ, Struck J, Morgenthaler NG, Bergmann A, Blankenbom S, Kee F, Nilsson P, Yin X, Peltonen L, Vartiainen E, Salomaa V, Hirschhorn JN, Melander O, Wang TJ. Association of common variants in *NPPA* and *NPPB* with circulating natriuretic peptides and blood pressure. *Nat Genet*. 2009;41(3):348-353.
  39. Clarke R, Peden JF, Hopewell JC, Kyriakou T, Goel A, Heath SC, Parish S, Barlera S, Franzosi MG, Rust S, Bennett D, Silveira A, Malarstig A, Green FR, Lathrop M, Gigante B, Leander K, de Faire U, Seedorf U, Hamsten A, Collins R, Watkins H, Farrall M; PROCARDIS Consortium. Genetic variants associated with Lp(a) lipoprotein level and coronary disease. *N Engl J Med*. 2009;361(26):2518-2528.
  40. Brunet G, Cerlich B, Robert P, Dumas S, Souetre E, Darcourt G. Open trial of a calcium antagonist, nimodipine, in acute mania. *Clin Neuropharmacol*. 1990;13(3):224-228.
  41. Pazzaglia PJ, Post RM, Ketter TA, Callahan AM, Marangell LB, Frye MA, George MS, Kimbrell TA, Leverich GS, Cora-Locatelli G, Luckenbaugh D. Nimodipine monotherapy and carbamazepine augmentation in patients with refractory recurrent affective illness. *J Clin Psychopharmacol*. 1998;18(5):404-413.